



**Manchester
Metropolitan
University**

Fatoye, Francis ORCID logoORCID: <https://orcid.org/0000-0002-3502-3953>, Mbada, Chidozie E, Gebrye, Tadesse ORCID logoORCID: <https://orcid.org/0000-0001-7976-2013>, Ogunsola, AO, Fatoye, Clara and Oyewole, O (2020) Real-World Prevalence of Musculoskeletal Disorders in Nigeria. In: ICHEP 2020 : 22th International Conference on Health Economics and Policy, 27 February 2020 - 28 February 2020, Sydney, Australia.

Downloaded from: <https://e-space.mmu.ac.uk/628883/>

Version: Accepted Version

Please cite the published version

<https://e-space.mmu.ac.uk>

Real-World Prevalence of Musculoskeletal Disorders in Nigeria

Authors : F. Fatoye, C. E. Mbada, T. Gebrye, A. O. Ogunsola, C. Fatoye, O. Oyewole

Abstract : Musculoskeletal disorders (MSDs) are a major cause of pain and disability. It is likely to become a greater economic and public health burden that is unnecessary. Thus, reliable prevalence figures are important for both clinicians and policy-makers to plan health care needs for those affected with the disease. This study estimated hospital based real-world prevalence of MSDs in Nigeria. A review of medical charts for adult patients attending Physiotherapy Outpatient Clinic at the Obafemi Awolowo University Teaching Hospitals Complex, Osun State, Nigeria between 2009 and 2018 was carried out to identify common MSDs including low back pain (LBP), cervical spondylosis (CSD), post immobilization stiffness (PIS), sprain, osteoarthritis (OA), and other conditions. Occupational class of the patients was determined using the International Labour Classification (ILO). Data were analysed using descriptive statistics of frequency and percentages. Overall, medical charts of 3,340 patients were reviewed within the span of ten years (2009 to 2018). Majority of the patients (62.8%) were in the middle class, and the remaining were in low class (25.1%) and high class (10.5%) category. An overall prevalence of 47.35% of MSD was found within the span of ten years. Of this, the prevalence of LBP, CSD, PIS, sprain, OA, and other conditions was 21.6%, 10%, 18.9%, 2%, 6.3%, and 41.3%, respectively. The highest (14.2%) and lowest (10.5%) prevalence of MSDs was recorded in the year of 2012 and 2018, respectively. The prevalence of MSDs is considerably high among Nigerian patients attending outpatient a physiotherapy clinic. The high prevalence of MSDs underscores the need for clinicians and decision makers to put in place appropriate strategies to reduce the prevalence of these conditions. In addition, they should plan and evaluate healthcare services to improve the health outcomes of patients with MSDs. Further studies are required to determine the economic burden of the condition and examine the clinical and cost-effectiveness of physiotherapy interventions for patients with MSDs.

Keywords : musculoskeletal disorders, Nigeria, prevalence, real world

Conference Title : ICHEP 2019 : International Conference on Health Economics and Policy

Conference Location : Sydney, Australia

Conference Dates : February 27-28, 2020